arXiv:1509.03400v1 [math.NT] 11 Sep 2015

NOTES ON WEIERSTRASS POINTS OF MODULAR CURVES $X_0(N)$

BO-HAE IM, DAEYEOL JEON AND CHANG HEON KIM

Abstract. We give conditions when the fixed points by the partial Atkin-Lehner involutions on $X_0(N)$ are Weierstrass points as an extension of the result by Lehner and Newman [16]. Furthermore, we complete their result by determining whether the fixed points by the full Atkin-Lehner involutions on $X_0(N)$ are Weierstrass points or not.

1. Introduction

Let $\mathfrak{H}$ be the complex upper half plane and Γ be a congruence subgroup of the full modular group $\mathrm{SL}_2(\mathbb{Z})$. Denote by $X(\Gamma)$ the modular curve obtained by compactifying the quotient $\Gamma\backslash\mathfrak{H}$. We can view $X(\Gamma)$ as a compact Riemann surface analytically. For each positive integer N, we let $\Gamma_0(N)$ be the Hecke subgroup of $\mathrm{SL}_2(\mathbb{Z})$ defined by

$$\Gamma_0(N) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z}) \,|\, c \equiv 0 \pmod{N} \right\}$$

and let $X_0(N) := X(\Gamma_0(N))$.

We recall the definitions of a Weierstrass point of a compact Riemann surface and its Weierstrass weight as in [2]. One says that a point P of a compact Riemann surface X of genus $g \geq 2$ is *a Weierstrass point* of X if there is a holomorphic differential ω (not identically zero) with a zero of order $\geq g$ at P. If $P \in X$ and $\omega_1, \omega_2, \ldots, \omega_g$ form a basis for the holomorphic differentials on X with the property that

$$0 = \mathrm{ord}_P(\omega_1) < \mathrm{ord}_P(\omega_2) < \cdots < \mathrm{ord}_P(\omega_g),$$

then *the Weierstrass weight of* P is the non-negative integer defined by

$$\mathrm{wt}(P) := \sum_{j=1}^{g} (\mathrm{ord}_P(\omega_j) - j + 1).$$

Note that $\mathrm{wt}(P) > 0$ if and only if P is a Weierstrass point of X.

The Weierstrass points of modular curves have been studied: Lehner and Newman [16] has given conditions when the cusp at infinity is a Weierstrass point of $X_0(N)$ for $N = 4n, 9n$ and Atkin [1] has given conditions for the case of $N = p^2 n$ where p is a prime ≥ 5. Also, Ogg [17], Kohnen [14, 15] and Kilger [13] have given some conditions when the cusp at infinity is not a Weierstrass point of $X_0(N)$ for certain N. And Ono [18] and Rohrlich [19] have investigated

---

Date: September 14, 2015.

2010 *Mathematics Subject Classification.* Primary 14H55, Secondary 11F06 and 11G18.

Key words and phrases. Weierstrass points, modular curves.

Bo-Hae Im was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (NRF-2014R1A1A2053748).

Weierstrass points of $X_0(p)$ for various primes p. Recently in [11] we investigated Weierstrass points on coverings of $X_0(N)$.

The Weierstrass points have been illustrated as an important class in number theory. For example, they have been used to determine the group structure of the automorphism groups of compact Riemann surfaces.

The main purpose of this paper is to give some conditions of being the Weierstrass points of $X_0(N)$. To describe our main result in this paper more precisely, we recall the definition of the Atkin-Lehner involution. We call a positive divisor Q of N with $\gcd(Q, N/Q) = 1$ an *exact divisor* of N and denote it by $Q \| N$. For each $Q \| N$, consider the matrices of the form $\begin{pmatrix} Qx & y \\ Nz & Qw \end{pmatrix}$ with $x, y, z, w \in \mathbb{Z}$ and determinant Q. Then such a matrix defines a unique involution on $X_0(N)$ which is called the *Atkin-Lehner involution* and denoted by W_Q. If $Q = N$, then W_Q is called the *full Atkin-Lehner involution,* and otherwise it is said to be a *partial Atkin-Lehner involution.* Sometimes we regard W_Q as a matrix.

Lehner and Newman [16] have shown that the fixed points by the full Atkin-Lehner involution on $X_0(N)$ are Weierstrass points except possibly for finitely many N which are listed in Lemma 3.3. In order to obtain the result, they have applied Schöneberg's Theorem ([20]). Our main result in this paper is to determine explicitly when the fixed points by the partial Atkin-Lehner involutions on $X_0(N)$ are Weierstrass points as an extension of the result by Lehner and Newman [16]. Also in this paper we give an algorithm to generate $\Gamma_0(N)$-inequivalent fixed points by the full Atkin-Lehner involution and we provide a computational method which can take care of the exceptional cases listed in Lemma 3.3 which are not covered in the Lehner and Newman [16].

This paper is organized as follows. In Section 2, we recall the formula ([8]) for the number of fixed points on $X_0(N)$ by the Atkin-Lehner involutions and explain algorithmically how to generate $\Gamma_0(N)$-inequivalent fixed points of $X_0(N)$ by the full Atkin-Lehner involution W_N. In Section 3, we give conditions when fixed points by the Atkin-Lehner involutions are Weierstrass points. We apply the formula ([8]) of the number of fixed points on $X_0(N)$ by the Atkin-Lehner involutions and Schöneberg's Theorem ([20]), and we give some new formulae for the number of fixed points when so-called *the elliptic condition* (Definition 3.9) is satisfied and apply them to obtain the conditions for Weierstrass points. In Section 4, we consider the exceptional cases which are not determined solely by Schöneberg's Theorem ([20]) and the formula given in [8], and we give a computational explanation how to determine whether they are Weierstrass points or not.

2. FIXED POINTS BY ATKIN-LEHNER INVOLUTIONS

Let $X_0^{+Q}(N)$ be the quotient space of $X_0(N)$ by W_Q. Let $g_0(N)$ and $g_0^{+Q}(N)$ be the genera of $X_0(N)$ and $X_0^{+Q}(N)$ respectively. Then $g_0^{+Q}(N)$ is computed by the Riemann-Hurwitz formula as follows:

$$g_0^{+Q}(N) = \frac{1}{4}(2g_0(N) + 2 - \nu(Q)),$$

where $\nu(Q) := \nu(Q; N)$ is the number of fixed points on $X_0(N)$ by W_Q. We recall the formula for $\nu(Q)$.

Proposition 2.1. ([8]) *For each $Q \| N$, $\nu(Q)$ is given by*

$$\begin{aligned}
\nu(Q) = &\left(\prod_{p|N/Q} c_1(p)\right) h(-4Q) \\
&+ \begin{cases} \left(\prod_{p|N/Q} c_2(p)\right) h(-Q), & \text{if } 4 \le Q \equiv 3 \pmod 4, \\ 0, & \text{otherwise} \end{cases} \\
&+ \begin{cases} \prod_{p|N/2} \left(1 + \left(\frac{-4}{p}\right)\right), & \text{if } Q = 2, \\ 0, & \text{otherwise} \end{cases} \\
&+ \begin{cases} \prod_{p|N/3} \left(1 + \left(\frac{-3}{p}\right)\right), & \text{if } Q = 3, \\ 0, & \text{otherwise} \end{cases} \\
&+ \begin{cases} \prod_{p^k|N/Q} \left(p^{[\frac{k}{2}]} + p^{[\frac{k-1}{2}]}\right), & \text{if } Q = 4, \\ 0, & \text{otherwise} \end{cases}
\end{aligned} \tag{2.1}$$

where $h(-d)$ is the class number of primitive quadratic forms of discriminant $-d$, $\left(\frac{\cdot}{\cdot}\right)$ is the Kronecker symbol and the functions $c_i(p)$ are defined as follows: for $i = 1, 2$,

$$c_i(p) = \begin{cases} 1 + \left(\frac{-Q}{p}\right), & \text{if } p \neq 2 \text{ and } Q \equiv 3 \pmod 4, \\ 1 + \left(\frac{-4Q}{p}\right), & \text{if } p \neq 2 \text{ and } Q \not\equiv 3 \pmod 4, \end{cases}$$

$$c_1(2) = \begin{cases} 1, & \text{if } Q \equiv 1 \pmod 4 \text{ and } 2\|N, \\ 0, & \text{if } Q \equiv 1 \pmod 4 \text{ and } 4|N, \\ 2, & \text{if } Q \equiv 3 \pmod 4 \text{ and } 2\|N, \\ 3 + \left(\frac{-Q}{2}\right), & \text{if } Q \equiv 3 \pmod 4 \text{ and } 4\|N, \\ 3\left(1 + \left(\frac{-Q}{2}\right)\right), & \text{if } Q \equiv 3 \pmod 4 \text{ and } 8|N, \end{cases}$$

$$c_2(2) = 1 + \left(\frac{-Q}{2}\right), \quad \text{if } Q \equiv 3 \pmod 4.$$

Next we give an algorithm to find $\Gamma_0(N)$-inequivalent fixed points of W_N on $X_0(N)$.

For a positive integer d conguent to 0 or 3 modulo 4, we denote by $\mathcal{Q}_d$ the set of positive definite integral binary quadratic forms

$$Q(x, y) = [a, b, c] := ax^2 + bxy + cy^2$$

with discriminant $-d = b^2 - 4ac$. Then $\mathrm{SL}_2(\mathbb{Z})$ acts on $\mathcal{Q}_d$ by

$$Q \circ \gamma(x, y) = Q(px + qy, rx + sy)$$

where $\gamma = \begin{pmatrix} p & q \\ r & s \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z})$. We say that a quadratic form $[a, b, c] \in \mathcal{Q}_d$ is *primitive* if $\gcd(a, b, c) = 1$. A primitive positive definite form $[a, b, c]$ is said to be *reduced* if

$$|b| \le a \le c, \text{ and } b \ge 0 \text{ if either } |b| = a \text{ or } a = c. \tag{2.2}$$

Let $\mathcal{Q}_d^\circ \subset \mathcal{Q}_d$ be the subset of primitive forms. Then $\mathrm{SL}_2(\mathbb{Z})$ also acts on $\mathcal{Q}_d^\circ$. As is well known, there is a 1-1 correspondence between the set of classes $\mathcal{Q}_d^\circ/\mathrm{SL}_2(\mathbb{Z})$ and the set of reduced forms.

Now for a fixed positive integer N we let d be a positive integer such that $-d$ is congruent to a square modulo $4N$. We choose an integer β with $-d \equiv \beta^2 \pmod{4N}$. Then we define

$$\mathcal{Q}_{d,N}^\circ = \{[aN, b, c] \in \mathcal{Q}_d \mid \gcd(a,b,c) = 1\}$$

and

$$\mathcal{Q}_{d,N,\beta}^\circ = \{[aN, b, c] \in \mathcal{Q}_{d,N}^\circ \mid \ b \equiv \beta \pmod{2N}\}.$$

Then $\Gamma_0(N)$ acts on both $\mathcal{Q}_{d,N}^\circ$ and $\mathcal{Q}_{d,N,\beta}^\circ$. For $Q = [aN, b, c] \in \mathcal{Q}_{d,N}^\circ$ we define $Q|W_N = [cN, -b, a]$. We observe that this defines an action of W_N on the set $\mathcal{Q}_{d,N}^\circ/\Gamma_0(N)$.

Assume that $N \geq 5$ and W_N fixes $\Gamma_0(N)\tau \in X_0(N)$ for some $\tau \in \mathfrak{H}$. This means that $\left(\begin{smallmatrix} p & q \\ r & s \end{smallmatrix}\right) W_N\tau = \tau$ for some $\left(\begin{smallmatrix} p & q \\ r & s \end{smallmatrix}\right) \in \Gamma_0(N)$. Thus τ satisfies a quadratic equation

$$NsX^2 - (r + qN)X + p = 0,$$

whose discriminant D is given by $D = (r + qN)^2 - 4psN = (r' - q)^2N^2 - 4N$ where $r = r'N$. Since $D \leq 0$, we come up with $(r' - q)^2N \leq 4$. Since we have assumed that $N \geq 5$, we must have $r' - q = 0$ and therefore $D = -4N$. Thus one has $D = 4q^2N^2 - 4psN = -4N$, which gives $\gcd(p, qN) = 1$ and hence $\gcd(s, -2qN, p) = 1$ or 2. If $\gcd(s, -2qN, p) = 2$, then both p and s should be even. It then follows from $ps - q^2N = 1$ that $N \equiv 3 \pmod 4$. Now we set $Q_\tau := [sN, -2qN, p]$. Then we see that

$$Q_\tau \in \mathcal{Q}_{4N,N,0}^\circ \ \text{ or } \ \frac{1}{2}Q_\tau \in \mathcal{Q}_{N,N,N}^\circ$$

where the latter case may happen only if $N \equiv 3 \pmod 4$. Let $\mathcal{F}_N$ be the set of the fixed points of W_N on $X_0(N)$ and let

$$\mathcal{G}_N := \begin{cases} \mathcal{Q}_{4N,N,0}^\circ/\Gamma_0(N) \cup \mathcal{Q}_{N,N,N}^\circ/\Gamma_0(N), & \text{if } N \equiv 3 \pmod 4 \\ \mathcal{Q}_{4N,N,0}^\circ/\Gamma_0(N), & \text{otherwise,} \end{cases}$$

where the union is disjoint. Now we define a map $\phi : \mathcal{F}_N \to \mathcal{G}_N$ by

$$\phi(\Gamma_0(N)\tau) = \begin{cases} Q_\tau, & \text{if } (s, -2qN, p) = 1 \\ \frac{1}{2}Q_\tau, & \text{if } (s, -2qN, p) = 2. \end{cases}$$

Then we can check easily ϕ is well-defined.

On the other hand, given a quadratic form $Q \in \mathcal{Q}_{4N,N,0}^\circ/\Gamma_0(N)$ or $Q \in \mathcal{Q}_{N,N,N}^\circ/\Gamma_0(N)$ we can find a fixed point $\Gamma_0(N)\tau \in X_0(N)$ such that $Q = \phi(\Gamma_0(N)\tau)$. Thus the map ϕ is surjective.

Now we deduce from [9, Proposition in p.505] that the natural maps

$$\begin{aligned} &\mathcal{Q}_{4N,N,0}^\circ/\Gamma_0(N) \to \mathcal{Q}_{4N}^\circ/\mathrm{SL}_2(\mathbb{Z}) \text{ and} \\ &\mathcal{Q}_{N,N,N}^\circ/\Gamma_0(N) \to \mathcal{Q}_{N}^\circ/\mathrm{SL}_2(\mathbb{Z}) \text{ (when } N \equiv 3 \pmod 4) \end{aligned} \tag{2.3}$$

are well-defined and bijective.

From [8], we have the following formula for the number of fixed points of W_N on $X_0(N)$:

$$\nu(N) = \#\mathcal{F}_N = \delta_N h(-4N),$$

where $h(-4N)$ denotes $\#\mathcal{Q}_{4N}^{\circ}/\mathrm{SL}_2(\mathbb{Z})$ and

$$\delta_N = \begin{cases} 2, & \text{if } N \equiv 7 \pmod 8 \\ \frac{4}{3}, & \text{if } N \equiv 3 \pmod 8 \text{ and } N > 3, \\ 1, & \text{otherwise.} \end{cases}$$

Since it is well-known that

$$h(-4N) = \begin{cases} h(-N), & \text{if } N \equiv 7 \pmod 8 \\ 3h(-N), & \text{if } N \equiv 3 \pmod 8 \text{ and } N > 3, \end{cases} \tag{2.4}$$

we have $\#\mathcal{F}_N = \#\mathcal{G}_N$, and hence ϕ is a bijection.

Now we will find $\Gamma_0(N)$-inequivalent fixed points in $\mathcal{F}_N$ by finding $\Gamma_0(N)$-inequivalent quadratic forms in $\mathcal{G}_N$ which can be obtained by pulling back the reduced forms in $\mathcal{Q}_{4N}^{\circ}/\mathrm{SL}_2(\mathbb{Z})$ and $\mathcal{Q}_{N}^{\circ}/\mathrm{SL}_2(\mathbb{Z})$ through the maps (2.3). Before providing an algorithm, we need the following lemma.

Lemma 2.2. *For a fixed positive integer N we let d be a positive integer such that $-d$ is congruent to a square modulo $4N$. We choose an integer β with $-d \equiv \beta^2 \pmod{4N}$. Then the following statements are true.*

1. *Given a primitive quadratic form $Q \in \mathcal{Q}_d^{\circ}$ there exists a quadratic form $[a, b, c]$ which is $\mathrm{SL}_2(\mathbb{Z})$-equivalent with Q and $\gcd(a, N) = 1$.*
2. *Let K be a solution to the linear congruence equation $2aX + b \equiv -\beta \pmod{2N}$ and set $[A, B, C] := [a, b, c] \circ \left(\begin{smallmatrix} K & -1 \\ 1 & 0 \end{smallmatrix}\right)$. Then $[A, B, C]$ belongs to $\mathcal{Q}_{d,N,\beta}$.*

Proof. (1) follows from [6, Lemma 2.3 and Lemma 2.25].
(2) First we note that b and β have the same parity since $-d = b^2 - 4ac$ and $-d \equiv \beta^2 \pmod{4N}$. Thus we obtain that $2 = \gcd(2a, 2N) \mid (-\beta - b)$, which guarantees that the linear congruence $2aX + b \equiv -\beta \pmod{2N}$ is solvable. Since $B = -2aK - b \equiv \beta \pmod{2N}$ and $C = a$, we must have

$$-4Aa = -4AC = -d - B^2 \equiv \beta^2 - B^2 \equiv 0 \pmod{4N},$$

which yields that $N|A$ since $\gcd(a, N) = 1$. Hence one has $[A, B, C] \in \mathcal{Q}_{d,N,\beta}$, as desired. □

Now we summarize the procedures explained in the above as the following algorithm.

Algorithm 2.3. *The following steps implement an algorithm to find $\Gamma_0(N)$-inequivalent fixed points of W_N:*
STEP 1. *Set $(d, \beta) = (4N, 0)$ or $(d, \beta) = (N, N)$ (when $N \equiv 3 \pmod 4$).*
STEP 2. *Starting from a reduced form Q^{red} satisfying (2.2) we first find a quadratic form $[a, b, c]$ which is $\mathrm{SL}_2(\mathbb{Z})$-equivalent with Q^{red} and $\gcd(a, N) = 1$.*
STEP 3. *Set $[A, B, C] := [a, b, c] \circ \left(\begin{smallmatrix} K & -1 \\ 1 & 0 \end{smallmatrix}\right)$ where K is a solution to the linear congruence equation$2aX + b \equiv -\beta \pmod{2N}$. Then $[A, B, C]$ belongs to $\mathcal{Q}_{d,N,\beta}$.*
STEP 4. *Let $\tau = \frac{-B+\sqrt{-d}}{2A}$. Then $\Gamma_0(N)\tau$ gives a fixed point of W_N.*

3. WEIERSTRASS POINT ON $X_0(N)$ ARISING FROM THE FIXED POINTS OF ATKIN-LEHNER INVOLUTIONS

First, we recall Schöneberg's Theorem [20] as follows.

Theorem 3.1. ([20]) *Let g be the genus of the modular curve $X(\Gamma)$ of a congruence subgroup Γ of $\mathrm{SL}_2(\mathbb{Z})$ and let M be an element of the normalizer of Γ in $\mathrm{SL}_2(\mathbb{R})$ and p be the exponent of M modulo Γ. Let g^* be the genus of the quotient space $X(\Gamma)/\langle M\rangle$ by the subgroup $\langle M\rangle$ generated by M modulo Γ. Then τ is a Weierstrass point on $X(\Gamma)$ provided that*

$$g^* \neq \lfloor g/p \rfloor.$$

($\lfloor x \rfloor$ denotes the largest integer not greater than x.)

By using the formula in Proposition 2.1 and Schöneberg's Theorem, we have the following:

Lemma 3.2. *Let τ be a fixed point of W_Q on $X_0(N)$ with $g_0(N) > 1$. If $\nu(Q) > 4$, then τ is a Weierstrass point of $X_0(N)$.*

Proof. By Theorem 3.1, τ is a Weierstrass point of $X_0(N)$ if

$$g_0(N) - 2g_0^{+Q}(N) > 1$$

which is equivalent to that $\nu(Q) > 4$. □

Note that $\nu(N) = \delta_N h(-4N)$ and there exist only finitely many N such that $h(-4N) \leq 4$. By using these facts, Lehner and Newman [16] have shown that the fixed points of W_N are Weierstrass points on $X_0(N)$ except possibly for finitely many N. However they didn't specify such possible N's, and hence we list them in Lemma 3.3. By Proposition 3.11 which we will prove later, we have the following:

Lemma 3.3. *The fixed points of W_N are Weierstrass points on $X_0(N)$ with $g_0(N) > 1$ except possibly for the following values for N:*

22, 28, 30, 33, 34, 37, 40, 42, 43, 45, 46, 48, 52, 57, 58, 60, 64, 67, 70, 72, 73, 78, 82, 85, 88, 93, 97, 100, 102, 112, 130, 133, 142, 148, 163, 177, 190, 193, 232, 253.

From now on, if $Q \| N$ and $N = QM$, we always assume $M > 1$. By Proposition 2.1 and Lemma 3.2 we have the following results.

Lemma 3.4. *Let $2 \| N$ and $N = 2M$. Then we have the following.*

1. *$\nu(2) = 0$ if and only if M has a prime factor p with $p \equiv 7 \pmod 8$ or prime factors q, r with $q \equiv 3 \pmod 8$ and $r \equiv 5 \pmod 8$.*
2. *If $\nu(2) \neq 0$, then*

$$\nu(2) = \delta_{0,s_2} 2^{s_0+s_1} + \delta_{0,s_1} 2^{s_0+s_2},$$

where s_0, s_1 and s_2 are the numbers of prime factors p of M with $p \equiv 1 \pmod 8$, $p \equiv 3 \pmod 8$ and $p \equiv 5 \pmod 8$ respectively, and $\delta_{i,j}$ is the Kronecker delta function.

Proof. Since $h(-8) = 1$,

$$\nu(2) = \prod_{p|M}\left(1 + \left(\frac{-8}{p}\right)\right) + \prod_{p|M}\left(1 + \left(\frac{-4}{p}\right)\right).$$

Since

$$\left(\frac{-8}{p}\right) = \begin{cases} 1, & p \equiv 1, 3 \pmod 8 \\ -1, & p \equiv 5, 7 \pmod 8 \end{cases}$$

and

$$\left(\frac{-4}{p}\right)=\begin{cases}1, & p\equiv 1 \pmod 4\\ -1, & p\equiv 3 \pmod 4\end{cases},$$

our result follows. □

Lemma 3.5. *Let $3\|N$ and $N=3M$. Then we have the following.*

1. $\nu(3)=0$ *if and only if* $8|M$ *or* M *has a prime factor* p *with* $p\equiv 5$ *or* $11 \pmod{12}$.
2. *If* $\nu(3)\neq 0$, *then*
$$\nu(3)=2^{s+1},$$
where s *is the number of prime factors* p *of* M *with* $p\equiv 1$ *or* $7 \pmod{12}$.

Proof. If $8|M$, then $\nu(3)=0$ because $\left(\frac{-3}{2}\right)=-1$. Suppose $8\nmid M$. Since $h(-12)=1$,

$$\nu(3)=\prod_{p|M}\left(1+(-1)^{p-1}\left(\frac{-3}{p}\right)\right)+\prod_{p|M}\left(1+\left(\frac{-3}{p}\right)\right)$$

Since

$$\left(\frac{-3}{p}\right)=\begin{cases}1, & p\equiv 1,7 \pmod{12}\\ -1, & p\equiv 5,11 \pmod{12}\end{cases}$$

for an odd prime p, the condition of $\nu(3)>0$ follows. If $\nu(3)>0$, then

$$\nu(3)=2\prod_{p|M}\left(1+\left(\frac{-3}{p}\right)\right)=2^{s+1}.$$

□

Lemma 3.6. *Let $4\|N$ and $N=4M$. Then W_4 has always a fixed point on $X_0(N)$. Let s and t denote the numbers of prime factors p of M with $p\equiv 1 \pmod 4$ and $p\equiv 3 \pmod 4$ respectively. Then we have the following:*

$$\nu(4)=\begin{cases}\prod_{p^k\|M}\left(p^{\left[\frac{k}{2}\right]}+p^{\left[\frac{k-1}{2}\right]}\right), & \text{if } t>0\\ \prod_{p^k\|M}\left(p^{\left[\frac{k}{2}\right]}+p^{\left[\frac{k-1}{2}\right]}\right)+2^s, & \text{if } t=0.\end{cases}$$

Proof. It follows directly from Proposition 2.1. □

By using lemmas 3.4, 3.5, and 3.6 we have the following result:

Theorem 3.7. *Let $Q\|N$ and $N=QM$. Suppose $\nu(Q)>0$. Then the fixed points of W_Q are Weierstrass points of $X_0(N)$ for each of the followings:*

1. $Q=2$; $s_0>1$ *and* $s_1=s_2=0$, $s_0+s_1>2$ *or* $s_0+s_2>2$ *where* s_0, s_1 *and* s_2 *are the numbers of prime factors* p,q *and* r *of* M *with* $p\equiv 1 \pmod 8$, $q\equiv 3 \pmod 8$ *and* $r\equiv 5 \pmod 8$ *respectively.*
2. $Q=3$; $s>1$ *where* s *is the number of prime factors* p *of* M *with* $p\equiv 1,7 \pmod{12}$.
3. $Q=4$; M *is not a square-free integer with* $M\neq 9$ *or* M *is square-free and* $6s+4t>11$ *where* s *and* t *are the numbers of prime factors* p *and* q *of* M *with* $p\equiv 1 \pmod 4$ *and* $q\equiv 3 \pmod 4$ *respectively.*

Proof. (1) and (2) follow immediately from Lemma 3.4 and Lemma 3.5.

For (3), if M is not a square-free integer with $M \neq 9$, then one can check easily that $\nu(4) > 4$ by Lemma 3.6. If M is square-free, then by Lemma 3.6

$$\nu(4) = \begin{cases} 2^{s+t}, & \text{if } t > 0 \\ 2^{s+1}, & \text{if } t = 0. \end{cases}$$

One can check easily that $\nu(4) > 4$ if and only if $6s + 4t > 11$. □

Now we deal with the case when $Q > 4$. For that we need the following result.

Proposition 3.8. *Let $Q \| N$ and $N = QM$. If $Q > 3$, then the following statements are equivalent.*

1. *W_Q has a fixed point on $X_0(N)$.*
2. *W_Q can be defined by a matrix of the form $\begin{pmatrix} Qx & y \\ Nz & -Qx \end{pmatrix}$ with $\det(W_Q) = Q$.*
3. *$X^2 \equiv -Q \pmod{N}$ has a solution.*
4. *$\left(\dfrac{-Q}{p}\right) = 1$ for any odd prime $p|M$, and $Q \equiv 3 \pmod 4$ if $4\|M$ and $Q \equiv 7 \pmod 8$ if $8|M$.*

Proof. (1)⇒(2) : Suppose $W_Q = \begin{pmatrix} Qx & y \\ Nz & Qw \end{pmatrix}$ with $\det(W_Q) = Q$ and W_Q has a fixed point on $X_0(N)$. Then there exists $\tau \in \mathfrak{H}$ such that $W_Q\tau = \gamma\tau$ for some $\gamma \in \Gamma_0(N)$. Since $\gamma^{-1}W_Q$ defines the same partial Atkin-Lehner involution, one may assume that $W_Q\tau = \tau$ by changing coefficients x, y, z, w of W_Q. Then the matrix W_Q is elliptic, and hence $|x + w|\sqrt{Q} < 2$. Since $Q > 3$, $x + w = 0$ and the result follows.

(2)⇒(3) : Suppose $W_Q = \begin{pmatrix} Qx & y \\ Nz & -Qx \end{pmatrix}$ with $\det(W_Q) = Q$. Then $\det(W_Q) = -(Qx)^2 - Nyz = Q$, and hence Qx is a solution of $X^2 \equiv -Q \pmod{N}$.

(3)⇒(2) : Suppose $X^2 \equiv -Q \pmod{N}$ has a solution, say x_0. Since $\gcd(Q, M) = 1$, one can choose Q' so that $QQ' \equiv 1 \pmod{M}$. Then $QQ'x_0$ is a solution of $X^2 \equiv -Q \pmod{N}$ which is divisible by Q. Letting $x = Q'x_0$, $(Qx)^2 = -Q - Nyz$ for some y, z, and hence the matrix $\begin{pmatrix} Qx & y \\ Nz & -Qx \end{pmatrix}$ has $\det(W_Q) = Q$.

(2)⇒(1) : Since $W_Q = \begin{pmatrix} Qx & y \\ Nz & -Qx \end{pmatrix}$ with $\det(W_Q) = Q$ is elliptic, it has a fixed point in $\mathfrak{H}$, and hence has one on $X_0(N)$.

(3)⇔(4) : It follows from [5, Ch.9, Theorem 9.13].

□

Definition 3.9. For $Q\|N$, if one of the equivalent statements of Proposition 3.8 is satisfied, then we call that $(N; Q)$ satisfies the *elliptic condition*.

From Proposition 2.1 and Proposition 3.8, we have the following:

Lemma 3.10. *Let $Q\|N$ and $N = QM$. Suppose $Q > 4$ and $(N;Q)$ satisfies the elliptic condition. Then the following holds: Let s be the number of prime divisors of M and*

$$\alpha_N = \begin{cases} 1, & \text{if } 2 \nmid N \text{ or } 2\|N, \\ 2, & \text{if } 4\|N, \\ 3, & \text{if } 8|N. \end{cases}$$

(1) *If $Q \equiv 7 \pmod 8$ or $Q \equiv 3 \pmod 8$ and N is odd, then*

$$\nu(Q) = 2^s(\alpha_N h(-4Q) + h(-Q)).$$

(2) *If $Q \equiv 1 \pmod 4$ and N is even, then*

$$\nu(Q) = 2^{s-1}h(-4Q).$$

(3) *If Q is even, or $Q \equiv 3 \pmod 8$ and N is even, or $Q \equiv 1 \pmod 4$ and N is odd, then*

$$\nu(Q) = 2^s h(-4Q).$$

Proof. If Q is even, then M is odd, and hence $c_1(p) = 2$ for all $p|M$. Thus $\nu(Q) = 2^s h(-4Q)$. This equality holds for the cases of $Q \equiv 1 \pmod 4$ and N is odd because M is odd. If $Q \equiv 1 \pmod 4$ and N is even, then M is even and $c_1(2) = 1$. Thus $\nu(Q) = 2^{s-1}h(-4Q)$. Consider the case $Q \equiv 3 \pmod 4$. If N is odd, then so is M, and $c_1(p) = c_2(p) = 2$ for all $p|M$. Thus $\nu(Q) = 2^s(h(-4Q)+h(-Q))$. If $Q \equiv 7 \pmod 8$ and N is even, then M is even, and $c_1(2) = 2\alpha_N$, $c_2(2) = c_i(p) = 2$ for all odd prime $p|M$ and $i = 1, 2$. Thus $\nu(Q) = 2^s(\alpha_N h(-4Q) + h(-Q))$. If $Q \equiv 3 \pmod 8$ and N is even, then M is even, and $c_2(2) = 0$, $c_1(p) = 2$ for all $p|M$. Thus $\nu(Q) = 2^s h(-4Q)$. □

For getting the condition that $\nu(Q) > 4$, we need to determine the values for Q with small $h(-Q)$ and $h(-4Q)$. Recall that if d_K is the (fundamental) discriminant of an imaginary quadratic field K, then $h(d_K)$ is equal to the class number of K, i.e. $h(d_K) = h(\mathcal{O}_K)$ where $\mathcal{O}_K$ is the ring of integers of K and $h(\mathcal{O}_K)$ is the order of the ideal class group of $\mathcal{O}_K$. On the other hand, if $d = f^2 d_K$ is the discriminant of a primitive quadratic form with $f > 1$, then $h(d) = h(\mathcal{O})$ where $\mathcal{O}$ is the order of conductor f in K (cf. [6]). Note that d is a fundamental discriminant if and only if one of the following statements holds:

- $d \equiv 1 \pmod 4$ and d is square-free,
- $d = 4m$ where $m \equiv 2$ or $3 \pmod 4$ and m is square-free.

The complete list of fundamental discriminants of class number 1 is as follows:

$$-3, -4, -7, -8, -11, -19, -43, -67, -163. \tag{3.1}$$

This is accomplished independently by Heegner [10], Baker [3] and Stark [21]. The non-fundamental discriminants of class number 1 are as follows:

$$-12, -16, -27, -28. \tag{3.2}$$

Thus $h(-Q) = 1$ if and only if $Q \in S_1$, where

$$S_1 = \{3, 4, 7, 8, 11, 12, 16, 19, 27, 28, 43, 67, 163\}.$$

The determination of fundamental discriminants of class number 2 was done again by Baker[4] and Stark[22].

Now consider the condition that $\nu(Q) > 4$. Due to (2.4) and Lemma 3.10, it suffices to determine the values for Q such that $h(-Q) = 1$ and $h(-4Q) = 2, 3, 4$. For the purpose, we refer to a paper by Klaise [12] in which all the orders of class number 2 and 3 are determined and an algorithm to find all orders of class number up to 100 is suggested. Let

$$\begin{aligned} S_2 &= \{5, 6, 8, 9, 10, 12, 13, 16, 18, 22, 25, 28, 37, 58\}, \\ S_3 &= \{11, 19, 23, 27, 31, 43, 67, 163\}, \\ S_4 &= \{14, 17, 20, 21, 24, 30, 32, 33, 34, 36, 39, 40, 42, 45, 46, 48, 49, \\ &\qquad 52, 55, 57, 60, 63, 64, 70, 72, 73, 78, 82, 85, 88, 93, 97, 100, 102, \\ &\qquad 112, 130, 133, 142, 148, 177, 190, 193, 232, 253\}. \end{aligned}$$

By using the result in [12] and some computations of the orders of class number 4 we have the following:

Proposition 3.11. *The list of Q with $h(-4Q) = 2, 3, 4$ is completely determined as follows:*

$$h(-4Q) = i \text{ if and only if } Q \in S_i \text{ for } i = 2, 3, 4.$$

By Proposition 3.11, we have the following result:

Theorem 3.12. *Let $Q \| N$ and $N = QM$. Suppose that $Q > 4$ and (N, Q) satisfies the elliptic condition. Let s denote the number of prime divisors of M.*

1. *When $Q \equiv 7 \pmod 8$ or $Q \equiv 3 \pmod 8$ and N is odd,*
   - a) *if $Q \neq 7$, then all the fixed points of W_Q on $X_0(N)$ are Weierstrass points, and*
   - b) *if $Q = 7$ and $4|N$ or $s > 1$, then all the fixed points of W_7 on $X_0(N)$ are Weierstrass points.*
2. *When $Q \equiv 1 \pmod 4$ and N is even,*
   - a) *if $Q \notin S_2 \cup S_4$, then all the fixed points of W_Q on $X_0(N)$ are Weierstrass points, and*
   - b) *if $Q \in S_2$ and $s > 2$ or $Q \in S_4$ and $s > 1$, then all the fixed points of W_Q on $X_0(N)$ are Weierstrass points.*
3. *In the other cases,*
   - a) *if $Q \notin S_2$, then all the fixed points of W_Q on $X_0(N)$ are Weierstrass points, and*
   - b) *if $Q \in S_2$ and $s > 1$, then all the fixed points of W_Q on $X_0(N)$ are Weierstrass points.*

Proof. (1) From Lemma 3.10, we need to consider Q such that $h(-4Q) + h(-Q) = 2$. By Proposition 3.11, one can check that the only $Q = 7$ such that $h(-4Q) + h(-Q) = 2$. In the case of $Q = 7$, if $4|N$ or $s > 1$, then $\nu(7) > 4$, and hence the result follows.

(2) From Lemma 3.10, we need to consider Q such that $h(-4Q) \leq 4$. By Proposition 3.11, one can check that there don't exist $Q \equiv 1 \pmod 4$ such that $h(-4Q) = 1, 3$, and hence the result follows.

(3) From Lemma 3.10, we need to consider Q such that $h(-4Q) \leq 2$. However by Proposition 3.11 one can check that there don't exist such a Q that $h(-4Q) = 1$ in these cases. Therefore the result follows. □

4. Computations for the exceptional cases

In this section, we completely determine when the fixed points of the full Atkin-Lehner involution W_N are Weierstrass points on $X_0(N)$ including the exceptional cases listed in Lemma 3.3. From now on, we always assume that N is one of those integer values. Let $\tau \in \mathfrak{H}$ so that $[\tau] := \Gamma_0(N)\tau \in X_0(N)$ is a fixed point of W_N. Let $\{f_1, f_2, \ldots, f_g\}$ be a basis of the cusp-form space $S_2(N)$ of weight 2 on $\Gamma_0(N)$, where g is the genus of $X_0(N)$. Then the differentials $\omega_i := f_i dz$ form a basis for the holomorphic differentials on $X_0(N)$. Since $\mathrm{ord}_{[\tau]}(\omega_i) = \mathrm{ord}_{[\tau]}(f_i)$, we will compute the Weierstrass weight of $[\tau]$ by using the f_i. In general, it is very difficult to compute the explicit forms of the f_i, and hence we will use the Fourier expansions at the infinite cusp ∞ of the f_i which can be easily computed by using the computer algebra system including Sage.

In general, the orders $\mathrm{ord}_{[\tau]}(f_i)$ do not satisfy the following:

$$0 = \mathrm{ord}_{[\tau]}(f_1) < \mathrm{ord}_{[\tau]}(f_2) < \cdots < \mathrm{ord}_{[\tau]}(f_g). \tag{4.1}$$

However, any basis satisfying (4.1) can be expressed as a linear combination of the f_i, and hence we have the following result:

Lemma 4.1. *Let $[\tau]$ be a fixed point of W_N on $X_0(N)$, and let $\{f_1, f_2, \ldots, f_g\}$ be a basis of the cusp form space $S_2(N)$ of weight 2 on $\Gamma_0(N)$. Then $[\tau]$ is a Weierstrass point of $X_0(N)$ if and only if $\det(A) = 0$, where $A = \left[\frac{f_i^{(j-1)}(\tau)}{(j-1)!}\right]$ is a $g \times g$ matrix with $1 \le i, j \le g$.*

Proof. First we consider the Taylor expansions of $f_i(z)$ at $z = \tau$ for $i = 1, \ldots, g$, which are given by

$$\begin{aligned}
f_1(z) &= f_1(\tau) + f_1'(\tau)(z-\tau) + \cdots + \frac{f_1^{(g-1)}(\tau)}{(g-1)!}(z-\tau)^{g-1} + \cdots \\
f_2(z) &= f_2(\tau) + f_2'(\tau)(z-\tau) + \cdots + \frac{f_2^{(g-1)}(\tau)}{(g-1)!}(z-\tau)^{g-1} + \cdots \\
&\ \ \vdots \\
f_g(z) &= f_g(\tau) + f_g'(\tau)(z-\tau) + \cdots + \frac{f_g^{(g-1)}(\tau)}{(g-1)!}(z-\tau)^{g-1} + \cdots .
\end{aligned}$$

By applying the elementary row operations to the above expansions we can find cusp forms $h_1, h_2, \ldots, h_g$ which are linear combinations of $f_1, f_2, \ldots, f_g$ with Taylor expansions of the form

$$\begin{aligned}
h_1(z) &= h_1(\tau) + h_1'(\tau)(z-\tau) + \cdots + \frac{f_1^{(g-1)}(\tau)}{(g-1)!}(z-\tau)^{g-1} + \cdots \\
h_2(z) &= \qquad\qquad h_2'(\tau)(z-\tau) + \cdots + \frac{h_2^{(g-1)}(\tau)}{(g-1)!}(z-\tau)^{g-1} + \cdots \\
&\qquad\qquad\qquad \ddots \\
h_g(z) &= \qquad\qquad\qquad\qquad\qquad\quad \frac{h_g^{(g-1)}(\tau)}{(g-1)!}(z-\tau)^{g-1} + \cdots
\end{aligned}$$

that satisfy

$$0 = \mathrm{ord}_P(\omega_1) < \mathrm{ord}_P(\omega_2) < \cdots < \mathrm{ord}_P(\omega_g),$$

where $P = [\tau]$ and $\omega_i = h_i dz$ for $i = 1, \ldots, g$. Thus we see that $\det(A) = c\prod_{j=1}^{g} \frac{h_j^{(j-1)}(\tau)}{(j-1)!}$ for some nonzero constant c and observe that

$$\begin{aligned} \det(A) = 0 &\iff \mathrm{ord}_P(\omega_j) \geq j \text{ for some } j \\ &\iff \mathrm{wt}(P) > 0 \\ &\iff P \text{ is a Weierstrass point.} \end{aligned}$$

□

Now we estimate the value $\frac{f_i^{(j-1)}(\tau)}{(j-1)!}$ by using the Fourier expansion of f_i at ∞ $f_i = \sum_{n=1}^{\infty} a_i(n)q^n$ where $z \in \mathfrak{H}$ and $q = e^{2\pi i z}$. Note that

$$\frac{f_i^{(j-1)}(\tau)}{(j-1)!} = \sum_{n=1}^{\infty} \frac{(2\pi i n)^{j-1}}{(j-1)!} a_i(n) q^n\big|_{z=\tau}. \tag{4.2}$$

Since $|e^{2\pi i\tau}| = e^{-2\pi \mathrm{Im}(\tau)}$, we need to find $\tau' \in [\tau]$ whose imaginary part is as big as possible for the fast convergence of the right side of (4.2). We will consider an algorithm to find $\tau' \in [\tau]$ so that $\mathrm{Im}(\tau')$ is the biggest. Note that $\mathrm{Im}(\alpha(\tau)) = \frac{\mathrm{Im}(\tau)}{|c\tau+d|}$ for $\alpha = \left(\begin{smallmatrix} a & b \\ c & d \end{smallmatrix}\right) \in \Gamma_0(N)$. Thus it suffices to find a pair (c, d) with $c \equiv 0 \pmod N$ and $\gcd(c, d) = 1$ so that $|c\tau + d|$ is the smallest. Let $\tau = x + iy$ with $x, y \in \mathbb{R}$. Since $|c\tau + d|^2 = (cx + d)^2 + (cy)^2 \leq 1$, there are only finitely many such pairs (c, d) because $|c| \leq \frac{1}{|y|}$ and $|cx + d| \leq 1$.

We summarize the procedures explained in the above as the following algorithm.

Algorithm 4.2. *The following steps implement an algorithm to find $\tau' \in [\tau]$ so that $\mathrm{Im}(\tau')$ is the biggest.*
STEP 1. *Set $m = [1/|y|]$ the largest integer not greater than $1/|y|$.*
STEP 2. *For each c with $-m \leq c \leq m$ and $c \equiv 0 \pmod N$, set $l_c = [-1-cx]$ and $u_c = [1-cx]$.*
STEP 3. *Pick a pair (c_0, d_0) so that $|c_0\tau+d_0|$ is the smallest among $|c\tau+d|$ where $-m \leq c \leq m$ and $l_c \leq d \leq u_c$ with $\gcd(c, d) = 1$.*
STEP 4. *Set $\tau' = \frac{\mathrm{Im}(\tau)}{|c_0\tau+d_0|}$.*

We list the fixed points of W_N in Table 1 by using Algorithm 2.3 and Algorithm 4.2. By using Lemma 4.1, we can confirm that there are no Weierstrass points on $X_0(N)$ arising from the fixed points of W_N. We have used Sage, Maple, and Mathematica for the numerical computations.

Table 1: Fixed points of W_N on $X_0(N)$

N	fixed points of W_N
22	$\frac{1}{\sqrt{-22}}, -\frac{6}{13} + \frac{\sqrt{-22}}{286}$
28	$\frac{1}{2\sqrt{-7}}, -\frac{8}{11} + \frac{\sqrt{-7}}{154}$

30	$\frac{1}{\sqrt{-30}}, \frac{8}{17} + \frac{\sqrt{-30}}{510}, \frac{4}{13} + \frac{\sqrt{-30}}{390}, \frac{2}{11} + \frac{\sqrt{-30}}{330}$
33	$\frac{1}{\sqrt{-33}}, \frac{1}{2} + \frac{\sqrt{-33}}{66}, \frac{9}{14} + \frac{\sqrt{-33}}{462}, -\frac{2}{7} + \frac{\sqrt{-33}}{231}$
34	$\frac{1}{\sqrt{-34}}, \frac{9}{19} + \frac{\sqrt{-34}}{646}, -\frac{4}{5} + \frac{\sqrt{-34}}{170}, \frac{4}{5} + \frac{\sqrt{-34}}{170}$
37	$\frac{1}{\sqrt{-37}}, \frac{1}{2} + \frac{\sqrt{-37}}{74}$
40	$\frac{1}{2\sqrt{-10}}, -\frac{6}{11} + \frac{\sqrt{-10}}{220}, -\frac{8}{13} + \frac{\sqrt{-10}}{260}, \frac{5}{7} + \frac{\sqrt{-10}}{140}$
42	$\frac{1}{\sqrt{-42}}, \frac{11}{23} + \frac{\sqrt{-42}}{966}, \frac{11}{17} + \frac{\sqrt{-42}}{714}, -\frac{2}{13} + \frac{\sqrt{-42}}{546}$
43	$\frac{1}{\sqrt{-43}}, -\frac{3}{4} + \frac{\sqrt{-43}}{172}, \frac{3}{4} + \frac{\sqrt{-43}}{172}, \frac{1}{2} + \frac{\sqrt{-43}}{86}$
45	$\frac{1}{3\sqrt{-5}}, \frac{1}{2} + \frac{\sqrt{-5}}{30}, \frac{11}{14} + \frac{\sqrt{-5}}{210}, \frac{4}{7} + \frac{\sqrt{-5}}{105}$
46	$\frac{1}{\sqrt{-46}}, \frac{12}{25} + \frac{\sqrt{-46}}{1150}, \frac{3}{5} + \frac{\sqrt{-46}}{230}, -\frac{3}{5} + \frac{\sqrt{-46}}{230}$
48	$\frac{1}{4\sqrt{-3}}, -\frac{13}{19} + \frac{\sqrt{-3}}{228}, -\frac{7}{13} + \frac{\sqrt{-3}}{156}, -\frac{1}{7} + \frac{\sqrt{-3}}{84}$
52	$\frac{1}{2\sqrt{-13}}, -\frac{13}{17} + \frac{\sqrt{-13}}{442}, \frac{4}{7} + \frac{\sqrt{-13}}{182}, -\frac{4}{7} + \frac{\sqrt{-13}}{182}$
57	$\frac{1}{\sqrt{-57}}, \frac{1}{2} + \frac{\sqrt{-57}}{114}, -\frac{15}{22} + \frac{\sqrt{-57}}{1254}, -\frac{4}{11} + \frac{\sqrt{-57}}{627}$
58	$\frac{1}{\sqrt{-58}}, \frac{15}{31} + \frac{\sqrt{-58}}{1798}$
60	$\frac{1}{2\sqrt{-15}}, \frac{15}{23} + \frac{\sqrt{-15}}{690}, \frac{14}{19} + \frac{\sqrt{-15}}{570}, \frac{10}{17} + \frac{\sqrt{-15}}{510}$
64	$\frac{1}{8\sqrt{-1}}, -\frac{9}{17} + \frac{\sqrt{-1}}{136}, -\frac{4}{5} + \frac{\sqrt{-1}}{40}, \frac{4}{5} + \frac{\sqrt{-1}}{40}$
67	$\frac{1}{\sqrt{-67}}, -\frac{3}{4} + \frac{\sqrt{-67}}{268}, \frac{3}{4} + \frac{\sqrt{-67}}{268}, \frac{1}{2} + \frac{\sqrt{-67}}{134}$
70	$\frac{1}{\sqrt{-70}}, \frac{18}{37} + \frac{\sqrt{-70}}{2590}, \frac{15}{19} + \frac{\sqrt{-70}}{1330}, \frac{12}{17} + \frac{\sqrt{-70}}{1190}$
72	$\frac{1}{6\sqrt{-2}}, \frac{9}{19} + \frac{\sqrt{-2}}{228}, -\frac{2}{17} + \frac{\sqrt{-2}}{204}, \frac{3}{11} + \frac{\sqrt{-2}}{132}$
73	$\frac{1}{\sqrt{-73}}, \frac{1}{2} + \frac{\sqrt{-73}}{146}, \frac{4}{7} + \frac{\sqrt{-73}}{511}, -\frac{4}{7} + \frac{\sqrt{-73}}{511}$
78	$\frac{1}{\sqrt{-78}}, \frac{20}{41} + \frac{\sqrt{-78}}{3198}, \frac{19}{29} + \frac{\sqrt{-78}}{2262}, -\frac{16}{19} + \frac{\sqrt{-78}}{1482}$
82	$\frac{1}{\sqrt{-82}}, \frac{21}{43} + \frac{\sqrt{-82}}{3526}, \frac{5}{7} + \frac{\sqrt{-82}}{574}, -\frac{5}{7} + \frac{\sqrt{-82}}{574}$
85	$\frac{1}{\sqrt{-85}}, \frac{1}{2} + \frac{\sqrt{-85}}{170}, \frac{13}{22} + \frac{\sqrt{-85}}{1870}, -\frac{2}{11} + \frac{\sqrt{-85}}{935}$
88	$\frac{1}{2\sqrt{-22}}, -\frac{12}{23} + \frac{\sqrt{-22}}{1012}, -\frac{12}{19} + \frac{\sqrt{-22}}{836}, -\frac{10}{13} + \frac{\sqrt{-22}}{572}$
93	$\frac{1}{\sqrt{-93}}, \frac{1}{2} + \frac{\sqrt{-93}}{186}, -\frac{23}{34} + \frac{\sqrt{-93}}{3162}, \frac{11}{17} + \frac{\sqrt{-93}}{1581}$
97	$\frac{1}{\sqrt{-97}}, \frac{1}{2} + \frac{\sqrt{-97}}{194}, -\frac{6}{7} + \frac{\sqrt{-97}}{679}, \frac{6}{7} + \frac{\sqrt{-97}}{679}$
100	$\frac{1}{10\sqrt{-1}}, -\frac{22}{29} + \frac{\sqrt{-1}}{290}, \frac{6}{13} + \frac{\sqrt{-1}}{130}, \frac{7}{13} + \frac{\sqrt{-1}}{130}$
102	$\frac{1}{\sqrt{-102}}, \frac{26}{53} + \frac{\sqrt{-102}}{5406}, -\frac{25}{37} + \frac{\sqrt{-102}}{3774}, \frac{19}{23} + \frac{\sqrt{-102}}{2346}$
112	$\frac{1}{4\sqrt{-7}}, -\frac{15}{29} + \frac{\sqrt{-7}}{812}, \frac{13}{23} + \frac{\sqrt{-7}}{644}, -\frac{7}{11} + \frac{\sqrt{-7}}{308}$
130	$\frac{1}{\sqrt{-130}}, \frac{33}{67} + \frac{\sqrt{-130}}{8710}, -\frac{25}{31} + \frac{\sqrt{-130}}{4030}, \frac{16}{23} + \frac{\sqrt{-130}}{2990}$
133	$\frac{1}{\sqrt{-133}}, \frac{1}{2} + \frac{\sqrt{-133}}{266}, -\frac{15}{26} + \frac{\sqrt{-133}}{3458}, \frac{11}{13} + \frac{\sqrt{-133}}{1729}$
142	$\frac{1}{\sqrt{-142}}, \frac{36}{73} + \frac{\sqrt{-142}}{10366}, -\frac{10}{11} + \frac{\sqrt{-142}}{1562}, \frac{10}{11} + \frac{\sqrt{-142}}{1562}$
148	$\frac{1}{2\sqrt{-37}}, -\frac{31}{41} + \frac{\sqrt{-37}}{3034}, -\frac{10}{19} + \frac{\sqrt{-37}}{1406}, \frac{10}{19} + \frac{\sqrt{-37}}{1406}$

163	$\frac{1}{\sqrt{-163}}, -\frac{3}{4} + \frac{\sqrt{-163}}{652}, \frac{3}{4} + \frac{\sqrt{-163}}{652}, \frac{1}{2} + \frac{\sqrt{-163}}{326}$
177	$\frac{1}{\sqrt{-177}}, \frac{1}{2} + \frac{\sqrt{-177}}{354}, \frac{41}{62} + \frac{\sqrt{-177}}{10974}, \frac{10}{31} + \frac{\sqrt{-177}}{5487}$
190	$\frac{1}{\sqrt{-190}}, \frac{48}{97} + \frac{\sqrt{-190}}{18430}, -\frac{26}{43} + \frac{\sqrt{-190}}{8170}, \frac{26}{29} + \frac{\sqrt{-190}}{5510}$
193	$\frac{1}{\sqrt{-193}}, \frac{1}{2} + \frac{\sqrt{-193}}{386}, -\frac{8}{11} + \frac{\sqrt{-193}}{2123}, \frac{8}{11} + \frac{\sqrt{-193}}{2123}$
232	$\frac{1}{2\sqrt{-58}}, -\frac{30}{59} + \frac{\sqrt{-58}}{6844}, \frac{23}{37} + \frac{\sqrt{-58}}{4292}, \frac{23}{31} + \frac{\sqrt{-58}}{3596}$
253	$\frac{1}{\sqrt{-253}}, \frac{1}{2} + \frac{\sqrt{-253}}{506}, -\frac{31}{34} + \frac{\sqrt{-253}}{8602}, -\frac{14}{17} + \frac{\sqrt{-253}}{4301}$

Acknowledgments. We would like to thank KIAS (Korea Institute for Advanced Study) for its hospitality while we have worked on this result. Daeyeol Jeon would like to thank Brown University for its hospitality during his sabbatical year.

REFERENCES

[1] A. O. L. Atkin, *Weierstrass points at cusps of* $\Gamma_0(n)$, Ann. Math., Second Series, **85** No.1 (Jan. 1967), 42–45.

[2] S. Ahlgren and K. Ono, *Weierstrass points on* $X_0(p)$ *and supersingular* j*-invariants*, Math. Ann., **325** (2003), 355–368.

[3] A. Baker, *Linear Forms in the logarithms of algebraic numbers I*, Mathematika, **13** (1996), 204–216.

[4] A. Baker, *Imaginary quadratic fields with class number 2*, Ann. Math., **94** (1971), 139–152.

[5] D. Burton, *Elementary Number Theory*, McGRAW-Hill, 2011.

[6] D. A. Cox, *Primes of the form* $x^2 + ny^2$*: Fermat, Class Field Theory, and Complex Multiplication*, John Wiley & Sons, Inc., New York, 1989.

[7] C. Delaunay, *Critical and ramification points of the modular parametrization of an elliptic curve*, J. Théor. Nombres Bordeaux **17** (2005), no. 1, 109-124.

[8] M. Furumoto and Y. Hasegawa, *Hyperelliptic quotients of modular curves* $X_0(N)$, Tokyo J. Math. **22** (1999), 105-125.

[9] Gross, B., Kohnen, W., Zagier, D.: *Heegner points and derivatives of L-series. II.* Math. Ann. **278**, 497–562 (1987).

[10] K. Heegner, *Diophantische Analysis und Modulfunktionen*, Math. Zeit. **56** (1952), 227-253.

[11] B.-H. Im, D. Jeon and C. H. Kim, *Weierstrass points on certain modular groups*, (2014), preprint.

[12] J. Klaise, *Orders in quadratic imaginary fields of small class number*, preprint.x‘x‘

[13] K. Kilger, *Weierstrass points on* $X_0(pl)$ *and arithmetic properties of Fourier coefficients of cusp forms*, Ramanujan J. **17** (2008), 321–330.

[14] W. Kohnen, *Weierstrass points at cusps on special modular curves*, Abh. Math. Sem. Univ. Hamburg **73** (2003), 241–251.

[15] W. Kohnen, *A short remark on Weierstrass points at infinity on* $X_0(N)$, Monatsh. Math. **143** (2004), 163–167.

[16] J. Lehner and M. Newman, *Weierstrass points of* $\Gamma_0(N)$, Ann of Math. (2) **79** (1964), 260–368.

[17] A. Ogg, *Weierstrass points of* $X_0(N)$, Illinois J. Math. **22** (1978), 31–35.

[18] K. Ono, *The web of modularity: arithmetic of the coefficients of modular forms and q-series*, CBMS Regional Conference Series in Mathematics **102**, Amer. Math. Soc., Providence, RI. 2004.

[19] D. Rohrlich, *Weierstrass points and modular forms*, Illinois J. Math. **29** (1985), 131-141.

[20] B. Schöneberg, *Über die Weierstrasspunkte in den Körpern der elliptischen Modulfunktionen*, Abh. Math. Sem. Univ. Hamburg **17** (1951), 104–111.

[21] H. M. Stark, *A complete determination of the complex quadratic fields of class number one*, Michigan Math. J., **14** (1967), 1–27.

[22] H. M. Stark, *On complex quadratic fields with class number two*, Math. Comp., **29** (1975), 289–302.

Bo-Hae Im, Department of Mathematics, Chung-Ang University, 84 Heukseok-Ro, Dongjak-Gu, Seoul 156-756, South Korea
E-mail address: `imbh@cau.ac.kr`

Daeyeol Jeon, Department of Mathematics education, Kongju National University, 56 Gongjudaehak-ro, Gongju-si, Chungcheongnam-do 314-701, South Korea
E-mail address: `dyjeon@kongju.ac.kr`

Chang Heon Kim, Department of Mathematics, Sungkyunkwan University, Suwon 440-746, South Korea
E-mail address: `chhkim@skku.edu`